

# *Gas and Stellar Properties of Simulated Galaxies in Cosmological Volume*

## *Gas Outflows*

**Paramita Bhattacharya**

(INAF - Astronomical Observatory of Trieste)

Trieste Numerical  
Cosmology Group



MUPPI collaborators:

Pierluigi Monaco,  
Giuseppe Murante,  
David Goz,  
Antonio Ragagnin,  
Matteo Viel



*Evolving Galaxies in Evolving Environments*

**Bologna**

**17 Sept. 2014**

# The Universe in a Box: Cosmological Hydrodynamic Simulations of Galaxy & Structure Formation

- Computational box  $\Leftrightarrow$  representative volume of the Universe
- Resolution elements (particles or grid) in box  $\Leftrightarrow$  matter
- Model LSS in terms of massive elements each of mass  $10^6 - 10^7 M_{\odot}$

## Steps:






### $\Lambda$ CDM cosmology



- Generate the initial condition
  - Primordial density fluctuations (Gaussian) at CMB epoch ( $z \sim 1100$ )
- Follow the non-linear evolution of density fields numerically
- Identify galaxies, clusters (group finder) at different  $z$
- Run on supercomputers
  - High speed & processing power, days to weeks of time



# Why Sub-Resolution Models ?



## Physics of baryons

- Radiative cooling and (photo + collisional) ionization heating of gas
- Fragmentation, clumping, multiphase ISM
- Star formation
- Metal production & chemical enrichment
- SN feedback, galactic wind
- AGN accretion + feedback
- ...


## In cosmological hydrodynamical simulations

(few - 10's Mpc) box : Resolution  $\sim 10^6 M_{Sun}$ , 1 kpc

- Baryonic physics, occurring on much smaller scales, are crucial ingredient
- Implemented as sub-resolution models

# Modified-GADGET<sub>3</sub> code: Numerical Sub-Resolution Physics

- GADGET : TreePM (gravity) - SPH (hydro)
  - Springel 2005, MNRAS, 364, 1105
- Metal-line cooling & radiative heating (Wiersma et al. 2009, MNRAS, 399, 574) in the presence of UV photoionizing background (Haardt & Madau 2001)
- Star Formation
- Stellar & Chemical Evolution (Tornatore et al. 2007, MNRAS, 382, 1050)
  - Metal (C, Ca, O, N, Ne, Mg, S, Si, Fe) release from SN type-II, type-Ia, & AGB stars; stellar age, mass & yield; different IMF; mass & metal loss from starburst
- SN Feedback
  - Thermal feedback ( $\uparrow T$ ) : inefficient, energy radiated away quickly
  - $\therefore$  Kinetic feedback ( $\uparrow v$ )
- No AGN feedback
  - In the simulations presented today



# Star-Formation in Multiphase ISM


- High-density SPH particle represents a part of ISM
  - Composed of 2 gas phases & stars

>  $SF, th$


COLD gas

STARS

HOT gas



- Effective model (*Springel & Hernquist 2003*)
  - Equilibrium solution
  - Self-regulated SF: constant effective pressure



- MUPPI = Multi-Phase Particle Integrator (*Murante et al. 2010*)
  - Molecular fraction of gas  $\propto$  Pressure
  - Mass & energy flows between components explicitly followed by numerically integrating system of ODEs within SPH timestep

# Existing Models of SN Feedback

- Kinetic feedback : give velocity kick to gas

- Energy-driven wind

- Springel & Hernquist (2003)

$$v_w = \text{constant}$$

$$v_w = 3\sigma_0 \sqrt{\frac{L}{L_{crit}} - 1}$$

Most of the models assume that wind velocity and mass-loading scales with some global galaxy property (mass, velocity dispersion, SFR)

- Radially-varying wind velocity

- Barai et al. (2013)

- Combinations & variations of energy and momentum-driven

- Schaye et al. (2010)
- Dave et al. (2013)
- Volgelberger et al. (2014)

- Thermal feedback : increase gas temperature


- Dalla Vecchia & Schaye (2012), Schaye et al. (2014)

- Turn off radiative cooling

- Stinson et al. (2006)


- ...

P. Barai, INAF-OATS



# SN Energy Feedback in MUPPI

- Energy imparted to gas particles
  - Inside SPH smoothing length and cone with semi-aperture angle =  $60^\circ$
  - Along path of least resistance
    - Negative density gradient
- Direct distribution of
  - Thermal energy
    - Efficiency fraction
    - Injected to local hot phase
  - Kinetic energy
    - Efficiency fraction, Probability
- No direct input expression of wind velocity & outflow mass loading




$$E_{th} = E_{SN} f_{fb,th} \frac{\Delta M_*}{M_{*,SN}}$$


$$E_{kin} = E_{SN} f_{fb,kin}$$

Large-scale filaments.  $(5/h \text{ Mpc})^3$  box at low-z. Dark matter - green, Gas - red, Stars - blue

Redshift: 0.170



Redshift: 0.170




Redshift: 0.170






# Simulation Runs (Barai et al. 2014, submitted)

Run Name	$L_{\text{box}}$ [Mpc]	$N_{\text{part}}$	$m_{\text{gas}}$ [ $M_{\odot}$ ]	$m_{\star}$ [ $M_{\odot}$ ]	$L_{\text{soft}}$ [kpc]	SF & SN feedback sub-resolution physics				
						Model	$v_w$	$f_{\text{fb,out}}$	$f_{\text{fb,kin}}$	$P_{\text{kin}}$
<i>E35nw</i>	35.56	$2 \times 320^3$	$8.72 \times 10^6$	$2.18 \times 10^6$	2.77 (comoving)	Effective	0			
<i>E35rvw</i>	35.56	$2 \times 320^3$	$8.72 \times 10^6$	$2.18 \times 10^6$	2.77 (comoving)	Effective	$v_w(r)$			
<i>E25cw</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	Effective	350			
<i>M25std</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.2	0.6	0.03
<i>M25a</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.4	0.4	0.03
<i>M25b</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.2	0.8	0.03
<i>M25c</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.2	0.6	0.01
<i>M25d</i>	25	$2 \times 256^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.2	0.6	0.06
<i>M50std</i>	50	$2 \times 512^3$	$5.36 \times 10^6$	$1.34 \times 10^6$	0.69 (physical)	MUPPI		0.2	0.5	0.03



# SFRD Evolution


# Outflow measurement technique (modified from Antonio Ragagnin 2013, Master thesis)




➤ Transform galaxy coordinates s.t. cold gas disk is rotating in X-Y plane

- Select gas particles:
  - lying inside either cylinder
  - moving at a high-velocity,  $|v_z| > V_{limit,outflow}$

- if  $(z \cdot v_z > 0) \Rightarrow$  Outflow
- if  $(z \cdot v_z < 0) \Rightarrow$  Inflow



# Setting the lower velocity threshold for outflow measurement




Fixed value:  
Reveals correlations.  
Adopt this for  
outflow velocity

P. Barai, INAF-OATs


Escape velocity:  
Measures escape outside halo  
Adopt this for  
mass outflow rate

# Outflow velocity vs. galaxy SFR




Observation : Martin (2005), Grimes et al. (2009), Banerji et al. (2011), Bordoloi et al. (2013)  
- positive correlation of outflow speed with galaxy mass and SFR.

# Mass outflow rate vs. galaxy SFR





# Mass loading factor ( $\eta = \text{Mass outflow rate} / \text{SFR}$ ) vs. halo mass






# Redshift Evolution of Outflow Number Fraction




Observation : Karman et al. (2014) - incidence of large-velocity outflow higher at  $z \sim 3$  than at  $z < 1$ .


# Redshift Evolution of Outflow Velocity vs SFR



# Redshift Evolution of Mass-Loading factor vs Halo Mass




# Realistic spiral galaxies in zoom-in cosmological simulations using moderate resolution (*Murante et al. 2014 submitted*)



**Figure 1.** Projected gas (upper panels) and stellar (lower panels) density for the GA2 simulation. The z-axis of the coordinate system is aligned with the angular momentum vector of the gas enclosed within the inner 8 kpc. Left panels show face-on densities, right column shows edge-on densities. Box size is 57 kpc.

# Study of bars in zoom-in cosmological simulations of spiral galaxies (Goz et al. 2014 submitted)



3-oct-14

# Summary

## SN feedback in cosmological hydrodynamic simulations:


- Can study impact of galactic winds on galaxy & IGM properties
  - Still far away from self-consistently driving these winds in such sims
  - Need subgrid prescription
- MUPPI is more physically-motivated sub-resolution model that uses only local properties of gas and generates realistic:
  - Galactic outflows
    - Outflow velocity positive correlation with global galaxy SFR
    - Constant mass-loading value at  $z=2$
    - Redshift evolution predicted over  $z = 1 - 5$
    - Need more observational data
  - Disk galaxies
- Future :
  - Compute further galaxy & IGM observables from sim
  - More physics : molecular cooling, AGN feedback

Formation of a disk galaxy at  $z=2$ .


Dark matter, gas, stars.

Edge-on view.


Redshift: 18.810



Redshift: 18.810






Redshift: 18.810



Formation of a disk galaxy at  $z=2$ .

Dark matter, gas, stars.

Face-on view.







# Extra Slides

# Mass outflow rate at $R_{\text{gal}}$ versus that at $R_{\text{vir}}$

Method	$N_{\text{outflow}}$	$f_{\text{outflow}}$
At $R_{\text{gal}}$ using $ v_r  > v_{\text{esc}}(R_{\text{gal}})$ , in a cylinder	1842	0.93
At $R_{\text{gal}}$ using $ v_r  > v_{\text{esc}}(R_{\text{gal}})$ , in a sphere	1936	0.97
At $R_{\text{vir}}$ using $ v_r  > v_{\text{esc}}(R_{\text{vir}})$ , in a sphere	1734	0.87



# Prediction with Theoretical Estimate of the MUPPI model

